

Die Size: 3" x 4"
Front

NataChew®

Prenatal Multivitamin
and Multimineral Tablets



Rx Only

DESCRIPTION:

Each tablet contains:

Supplement Facts

Serving Size: 1 Tablet

	Amount Per Serving
Vitamin A (beta carotene)	2,700 IU
Vitamin C (sodium ascorbate and ascorbic acid)	120 mg
Vitamin D ₃ (cholecalciferol)	400 IU
Vitamin E (d-alpha tocopheryl acetate)	20 IU
Vitamin B ₁ (thiamine mononitrate)	2 mg
Vitamin B ₂ (riboflavin)	3 mg
Vitamin B ₆ (pyridoxine HCl)	10 mg
Folic Acid	1 mg
Vitamin B ₁₂ (cyanocobalamin)	12 mcg
Iron (ferrous fumarate and ferrous II bis-glycinate chelate)	28 mg
Niacinamide	20 mg

OTHER INGREDIENTS: sugar, D&C red #7 calcium lake, sucralose, flavor, microcrystalline cellulose, polyethylene glycol 400, stearic acid, magnesium stearate

INDICATIONS AND USAGE: NataChew® is a prescription pre/post-natal multivitamin, multimineral indicated for use in improving the nutritional status of women throughout pregnancy, as well as in the post-natal period for both lactating and non-lactating mothers. This product is also useful in improving the nutritional status of women prior to conception.

CONTRAINDICATIONS: This product is contraindicated in patients with known hypersensitivity to any of the ingredients.

WARNINGS AND PRECAUTIONS:

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6. Keep this product out of reach of children. In case of accidental overdose, call a doctor or poison control center immediately.

Folic acid is improper therapy in the treatment of pernicious anemia in that hematological remission can occur while neurological manifestations remain progressive. While prescribing this product for pregnant women, nursing mothers, or for women prior to conception, their medical condition and use of other drugs, herbs, and/or supplements should be considered.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

ADVERSE REACTIONS: Allergic sensitization has been reported following both oral and parenteral administration of folic acid. **You should call your doctor for medical advice about side effects. To report a serious adverse event, call 1-855-899-4237.**

DOSAGE AND ADMINISTRATION: One tablet daily with or without food or as prescribed by a physician.

HOW SUPPLIED: NataChew® is supplied as a red tablet, imprinted "902", in child-resistant bottles of 90 tablets, NDC 44118-902-90, and in sample bottles of 5 tablets, NDC 44118-902-05.

Store between 15°C and 30°C (between 59°F and 86°F). Tamper evident by foil seal under cap.

Manufactured for:
Eckson Labs
1000 N. West St., Suite 1200, #927
Wilmington, DE 19801
vAT Rev. 09/2012



902-051-02